



ROGER L. BLAKE, EXECUTIVE DIRECTOR

CALIFORNIA INTERSCHOLASTIC FEDERATION

CIF STATE OFFICE • 4658 DUCKHORN DRIVE • SACRAMENTO, CA 95834 • PH: 916-239-4477 • FX: 916-239-4478 • WWW.CIFSTATE.ORG

Bylaw 503 H. Concussion Protocols as Per Ed. Code 49475.(a).(1)

A student-athlete who is suspected of sustaining a concussion or head injury in a practice or game shall be removed from competition at that time for the remainder of the day. A student-athlete who has been removed from play may not return to play until the athlete is evaluated by a licensed health care provider trained in education and management of concussion and receives written clearance to return to play from that health care provider. If a licensed health care provider, trained in education and management of concussion determines that the athlete sustained a concussion or a head injury, the athlete is required to complete a graduated return-to-play protocol of no less than 7 days, from the time of diagnosis, in duration under the supervision of a licensed health care provider. On a yearly basis, a concussion and head injury information sheet shall be signed and returned by all athletes and the athlete's parent or guardian before the athlete's initiating practice or competition.

PARENT & ATHLETE CONCUSSION INFORMATION SHEET



WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by a bump, blow, or jolt to the head or body that causes the head and brain to move quickly back and forth. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports one or more symptoms of concussion after a bump, blow, or jolt to the head or body, s/he should be kept out of play the day of the injury. The athlete should only return to play with permission from a health care professional experienced in evaluating for concussion.

DID YOU KNOW?

- Most concussions occur without loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.



SYMPTOMS REPORTED BY ATHLETE:

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not “feeling right” or is “feeling down”

SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can’t recall events prior to hit or fall
- Can’t recall events after hit or fall



“IT’S BETTER TO MISS ONE GAME THAN THE WHOLE SEASON”

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

1. If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.
2. Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, and playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.
3. Remember: Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. They can even be fatal.

STUDENT-ATHLETE NAME PRINTED

STUDENT-ATHLETE NAME SIGNED

DATE

PARENT OR GUARDIAN NAME PRINTED

PARENT OR GUARDIAN NAME SIGNED

DATE

JOIN THE CONVERSATION  www.facebook.com/CDCHeadsUp



HEADS UP

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

CONCUSSION FACT SHEET FOR COACHES



WHAT IS A CONCUSSION?

Concussion, a type of traumatic brain injury, is caused by a bump, blow, or jolt to the head. Concussions can also occur from a blow to the body that causes the head and brain to move rapidly back and forth—literally causing the brain to bounce around or twist within the skull.

This sudden movement of the brain causes stretching and tearing of brain cells, damaging the cells and creating chemical changes in the brain.

HOW CAN I RECOGNIZE A POSSIBLE CONCUSSION?

Concussions can result from a fall or from athletes colliding with each other, the ground, or with an obstacle, such as a goalpost. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

As a coach you are on the front line in identifying an athlete with a suspected concussion. You know your athletes well and can recognize when something is off—even when the athlete doesn’t know it or doesn’t want to admit it.

So to help spot a concussion, you should watch for and ask others to report the following two things:

1. A forceful bump, blow, or jolt to the head or body that results in rapid movement of the head.

AND

2. Any concussion signs or symptoms, such as a change in the athlete’s behavior, thinking, or physical functioning.

Signs and symptoms of concussion generally show up soon after the injury. But the full effect of the injury may not be noticeable at first. For example, in the first few minutes the athlete might be slightly confused or appear a little bit dazed, but an hour later they can’t recall coming to the practice or game.

You should repeatedly check for signs of concussion and also tell parents what to watch out for at home. Any worsening of concussion signs or symptoms indicates a medical emergency.

SIGNS AND SYMPTOMS

Athletes who experience one or more of the signs and symptoms listed below, or who report that they just “don’t feel right,” after a bump, blow, or jolt to the head or body, may have a concussion.

SYMPTOMS REPORTED BY ATHLETE:

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not “feeling right” or is “feeling down”

SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can’t recall events prior to hit or fall
- Can’t recall events after hit or fall

WHAT ARE CONCUSSION DANGER SIGNS?

In rare cases, a dangerous blood clot may form on the brain in an athlete with a concussion and crowd the brain against the skull. Call 9-1-1 or take the athlete to the emergency department right away if after a bump, blow, or jolt to the head or body the athlete exhibits one or more of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

FACTS

Sometimes people wrongly believe that it shows strength and courage to play injured. Some athletes may also try to hide their symptoms.

Don't let your athlete convince you that he or she is "just fine" or that he or she can "tough it out." Discourage others from pressuring injured athletes to play. Emphasize to athletes and parents that playing with a concussion is dangerous.

WHAT SHOULD I DO IF A CONCUSSION IS SUSPECTED?

No matter whether the athlete is a key member of the team or the game is about to end, an athlete with a suspected concussion should be immediately removed from play. To help you know how to respond, follow the Heads Up four-step action plan:

1. REMOVE THE ATHLETE FROM PLAY.

Look for signs and symptoms of a concussion if your athlete has experienced a bump or blow to the head or body. When in doubt, sit them out!

2. ENSURE THAT THE ATHLETE IS EVALUATED BY AN APPROPRIATE HEALTH CARE PROFESSIONAL.

Do not try to judge the severity of the injury yourself. Health care professionals have a number of methods that they can use to assess the severity of concussions. As a coach, recording the following information can help health care professionals in assessing the athlete after the injury:

- Cause of the injury and force of the hit or blow to the head or body
- Any loss of consciousness (passed out/knocked out) and if so, for how long
- Any memory loss immediately following the injury
- Any seizures immediately following the injury
- Number of previous concussions (if any)

3. INFORM THE ATHLETE'S PARENTS OR GUARDIANS.

Let them know about the possible concussion and give them the Heads Up fact sheet for parents. This fact sheet can help parents monitor the athlete for sign or symptoms that appear or get worse once the athlete is at home or returns to school.

4. KEEP THE ATHLETE OUT OF PLAY.

An athlete should be removed from play the day of the injury and until an appropriate health care professional says they are symptom-free and it's OK to return to play. After you remove an athlete with a suspected concussion from practice or play, the decision about return to practice or play is a medical decision.



WHY SHOULD I BE CONCERNED ABOUT CONCUSSIONS?

Most athletes with a concussion will recover quickly and fully. But for some athletes, signs and symptoms of concussion can last for days, weeks, or longer.

If an athlete has a concussion, his or her brain needs time to heal. A repeat concussion that occurs before the brain recovers from the first—usually within a short time period (hours, days, weeks)—can slow recovery or increase the chances for long-term problems. In rare cases, repeat concussions can result in brain swelling or permanent brain damage. It can even be fatal.



DID YOU KNOW?

- Young children and teens are more likely to get a concussion and take longer to recover than adults.
- Athletes who have ever had a concussion are at increased risk for another concussion.
- All concussions are serious.
- Recognition and proper response to concussions when they first occur can help prevent further injury or even death.

HOW CAN I HELP ATHLETES TO RETURN TO PLAY GRADUALLY?

An athlete should return to sports practices under the supervision of an appropriate health care professional. When available, be sure to work closely with your team's certified athletic trainer.

Below are five gradual steps that you and the health care professional should follow to help safely return an athlete to play. Remember, this is a gradual process. These steps should not be completed in one day, but instead over days, weeks, or months.

BASELINE:

Athletes should not have any concussion symptoms. Athletes should only progress to the next level of exertion if they do not have any symptoms at the current step.

STEP 1:

Begin with light aerobic exercise only to increase an athlete's heart rate. This means about 5 to 10 minutes on an exercise bike, walking, or light jogging. No weight lifting at this point.

STEP 2:

Continue with activities to increase an athlete's heart rate with body or head movement. This includes moderate jogging, brief running, moderate-intensity stationary biking, moderate-intensity weightlifting (reduced time and/or reduced weight from your typical routine).

STEP 3:

Add heavy non-contact physical activity, such as sprinting/running, high-intensity stationary biking, regular weightlifting routine, non-contact sport-specific drills (in 3 planes of movement).

STEP 4:

Athlete may return to practice and full contact (if appropriate for the sport) in controlled practice.

STEP 5:

Athlete may return to competition.

If an athlete's symptoms come back or she or he gets new symptoms when becoming more active at any step, this is a sign that the athlete is pushing him or herself too hard. The athlete should stop these activities and the athlete's health care provider should be contacted. After more rest and no concussion symptoms, the athlete should begin at the previous step.

HOW CAN I HELP PREVENT CONCUSSIONS OR OTHER SERIOUS BRAIN INJURIES?

Insist that safety comes first. To help minimize the risks for concussion or other serious brain injuries:

- Ensure that athletes follow the rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.
- Make sure the athlete wears the right protective equipment for their activity. Protective equipment should fit properly, be well maintained, and be worn consistently and correctly.
- Wearing a helmet is a must to reduce the risk of severe brain injury and skull fracture. However, a helmet doesn't make an athlete immune to concussion. There is no "concussion-proof" helmet.

Check with your league, school, or district about concussion policies. Concussion policy statements can be developed to include:

- The school or league's commitment to safety
- A brief description of concussion
- Information on when athletes can safely return to school and play.

Parents and athletes should sign the concussion policy statement at the beginning of the season.



▶ **"WHEN IN DOUBT,
SIT THEM OUT!"**




JOIN THE CONVERSATION AT www.facebook.com/CDCHeadsUp

TO LEARN MORE GO TO [WWW.CDC.GOV/CONCUSSION](http://www.cdc.gov/concussion)

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

HEADS UP

HEADS UP CONCUSSION ACTION PLAN



IF YOU SUSPECT THAT AN ATHLETE HAS A CONCUSSION, YOU SHOULD TAKE THE FOLLOWING STEPS:

1. Remove the athlete from play.
2. Ensure that the athlete is evaluated by a health care professional experienced in evaluating for concussion. Do not try to judge the seriousness of the injury yourself.
3. Inform the athlete's parents or guardians about the possible concussion and give them the fact sheet on concussion.
4. Keep the athlete out of play the day of the injury. An athlete should only return to play with permission from a health care professional, who is experienced in evaluating for concussion.

▶ **"IT'S BETTER TO MISS ONE GAME, THAN THE WHOLE SEASON."**



CONCUSSION SIGNS AND SYMPTOMS

Athletes who experience one or more of the signs and symptoms listed below after a bump, blow, or jolt to the head or body may have a concussion.

SYMPTOMS REPORTED BY ATHLETE

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

SIGNS OBSERVED BY COACHING STAFF

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can't recall events prior to hit or fall

JOIN THE CONVERSATION AT www.facebook.com/CDCHeadsUp

HEADS UP

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

CONCUSSION FACT SHEET FOR PARENTS



WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury. Concussions are caused by a bump or blow to the head. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

You can’t see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If your child reports any symptoms of concussion, or if you notice the symptoms yourself, seek medical attention right away.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

If your child has experienced a bump or blow to the head during a game or practice, look for any of the following signs of a concussion:

SYMPTOMS REPORTED BY ATHLETE:

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not “feeling right” or is “feeling down”

SIGNS OBSERVED BY PARENTS/ GUARDIANS:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes



DANGER SIGNS

Be alert for symptoms that worsen over time. Your child or teen should be seen in an emergency department right away if s/he has:

- One pupil (the black part in the middle of the eye) larger than the other
- Drowsiness or cannot be awakened
- A headache that gets worse and does not go away
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Difficulty recognizing people or places
- Increasing confusion, restlessness, or agitation
- Unusual behavior
- Loss of consciousness (even a brief loss of consciousness should be taken seriously)

WHAT SHOULD YOU DO IF YOU THINK YOUR CHILD HAS A CONCUSSION?

1. **SEEK MEDICAL ATTENTION RIGHT AWAY**
A health care professional will be able to decide how serious the concussion is and when it is safe for your child to return to regular activities, including sports.
2. **KEEP YOUR CHILD OUT OF PLAY.**
Concussions take time to heal. Don't let your child return to play the day of the injury and until a health care professional says it's OK. Children who return to play too soon - while the brain is still healing - risk a greater chance of having a second concussion. Repeat or later concussions can be very serious. They can cause permanent brain damage, affecting your child for a lifetime.
3. **TELL YOUR CHILD'S COACH ABOUT ANY PREVIOUS CONCUSSION.**
Coaches should know if your child had a previous concussion. Your child's coach may not know about a concussion your child received in another sport or activity unless you tell the coach.

HOW CAN YOU HELP YOUR CHILD PREVENT A CONCUSSION OR OTHER SERIOUS BRAIN INJURY?

- Ensure that they follow their coach's rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.
- Make sure they wear the right protective equipment for their activity. Protective equipment should fit properly and be well maintained.
- Wearing a helmet is a must to reduce the risk of a serious brain injury or skull fracture.
 - However, helmets are not designed to prevent concussions. There is no "concussion-proof" helmet. So, even with a helmet, it is important for kids and teens to avoid hits to the head.

HOW CAN I HELP MY CHILD RETURN TO SCHOOL SAFELY AFTER A CONCUSSION?

Children and teens who return to school after a concussion may need to:

- Take rest breaks as needed
- Spend fewer hours at school
- Be given more time to take tests or complete assignments
- Receive help with schoolwork
- Reduce time spent reading, writing, or on the computer

Talk with your child's teachers, school nurse, coach, speech-language pathologist, or counselor about your child's concussion and symptoms. As your child's symptoms decrease, the extra help or support can be removed gradually.



JOIN THE CONVERSATION  www.facebook.com/CDCHeadsUp

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

CONCUSSION FACT SHEET FOR ATHLETES



CONCUSSION FACTS

- A concussion is a brain injury that affects how your brain works.
- A concussion is caused by a bump, blow, or jolt to the head or body.
- A concussion can happen even if you haven't been knocked out.
- If you think you have a concussion, you should not return to play on the day of the injury and until a health care professional says you are OK to return to play.



CONCUSSION SIGNS AND SYMPTOMS

Concussion symptoms differ with each person and with each injury, and may not be noticeable for hours or days.

Common symptoms include:

- Headache
- Confusion
- Difficulty remembering or paying attention
- Balance problems or dizziness
- Feeling sluggish, hazy, foggy, or groggy
- Feeling irritable, more emotional, or "down"
- Nausea or vomiting
- Bothered by light or noise
- Double or blurry vision
- Slowed reaction time
- Sleep problems
- Loss of consciousness

During recovery, exercising or activities that involve a lot of concentration (such as studying, working on the computer, or playing video games) may cause concussion symptoms to reappear or get worse.

WHY SHOULD I REPORT MY SYMPTOMS?

- Unlike with some other injuries, playing or practicing with concussion symptoms is dangerous and can lead to a longer recovery and a delay in your return to play.
- While your brain is still healing, you are much more likely to have another concussion.
- A repeat concussion in a young athlete can result in permanent damage to your brain. They can even be fatal.

WHAT SHOULD I DO IF I THINK I HAVE A CONCUSSION?

DON'T HIDE IT. REPORT IT.

Ignoring your symptoms and trying to "tough it out" often makes symptoms worse. Tell your coach, parent, and athletic trainer if you think you or one of your teammates may have a concussion. Don't let anyone pressure you into continuing to practice or play with a concussion.

GET CHECKED OUT.

Only a health care professional can tell if you have a concussion and when it's OK to return to play. Sports have injury timeouts and player substitutions so that you can get checked out and the team can perform at its best. The sooner you get checked out, the sooner you may be able to safely return to play.

TAKE CARE OF YOUR BRAIN.

A concussion can affect your ability to do schoolwork and other activities. Most athletes with a concussion get better and return to sports, but it is important to rest and give your brain time to heal. A repeat concussion that occurs while your brain is still healing can cause long-term problems that may change your life forever.



▶ **“IT’S BETTER TO MISS ONE GAME,
THAN THE WHOLE SEASON.”**

JOIN THE CONVERSATION AT www.facebook.com/CDCHeadsUp



HEADS UP

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).